



ELSEVIER

Nuclear Instruments and Methods in Physics Research B xxx (2002) xxx–xxx

**NIM B**  
Beam Interactions  
with Materials & Atoms

www.elsevier.com/locate/nimb

## Can we measure the gravitational free fall of cold Rydberg state positronium?

A.P. Mills Jr.<sup>a,\*</sup>, M. Leventhal<sup>b</sup>

<sup>a</sup> Department of Physics, University of California, Riverside, CA 92521, USA

<sup>b</sup> Department of Astronomy, University of Maryland, College Park, MD, USA

### Abstract

In this paper we examine the possibilities for detecting the free fall of Rydberg positronium atoms. In our scheme, cold positronium atoms are emitted from a “point” source and excited to the  $n = 25$  circular Rydberg state with  $L = n - 1$ . The positronium atoms are allowed to travel horizontally 10 m in a field free vacuum and focused onto a detector using an elliptical Van der Waals mirror. A free fall distance of order 50  $\mu\text{m}$  and a few detected atoms per hour are anticipated. Various extraneous influences on the positronium, such as collisions with residual gas atoms, Stark mixing in stray electric and magnetic fields, photoionization due to thermal radiation, and accelerations due to patch potentials are estimated. © 2002 Published by Elsevier Science B.V.

### 1. Introduction

Notwithstanding the conventional point of view that antigravity is impossible within the established framework of modern physics, the hypothesis that antimatter is attracted to matter has not been the subject of a direct test. Attempts by Fairbank [1] and his group to measure the gravitational free fall of electrons and positrons were not conclusive and work is in now progress to prepare antihydrogen atoms for free fall measurements in vacuum [2]. A measurement of the free fall of neutral positronium atoms would be much simpler than for its constituent charged particles. However, the positronium must be in a high Rydberg state in order to have a sufficiently

long lifetime to result in an observable free fall distance. In such a state the positronium is highly polarizable and the free fall region must be nearly as free of stray electric and magnetic fields as would have been needed for observations on free electrons and positrons. Nevertheless, the experiment appears feasible with current technology.

The proposal is to produce a “point” source of positronium [3] of 1- $\mu\text{m}$  vertical extent and to excite it via Doppler-free techniques to an  $n \approx 25$  state. There are now several possibilities for a free fall experiment. The simplest perhaps is to allow the positronium in the low energy tail of its energy distribution to travel a few meters in the horizontal direction, be focused by a mirror and then travel a few more meters to converge on a 1- $\mu\text{m}$  spot on a position sensitive detector, as shown in Fig. 1. Deflections of the spot as large as 25  $\mu\text{m}$  should be observable at the longest positronium time of flight delay.

\* Corresponding author.

E-mail addresses: apmjr@citrus.ucr.edu (A.P. Mills Jr.), leventhal@astro.umd.edu (M. Leventhal).




Fig. 1. Experiment to measure the gravitational free fall of  $n = 25$  Rydberg positronium.

51 A second possibility using lasers to make an  
 52 atom interferometer and a third to use transmis-  
 53 sion gratings to make a Mach-Zender interfer-  
 54 ometer [4] need to be evaluated and compared to  
 55 the mirror experiment.

## 56 2. Cold positronium production

57 Positrons from the final brightness enhancement  
 58 remoderator of an intense slow positron beam  
 59 source are accelerated to 3 keV and brought to a  
 60 line focus of vertical width equal to  $1 \mu\text{m}$  and  
 61 horizontal width  $300 \mu\text{m}$  on an aluminum sample.  
 62 The sample is a single crystal Al(111) surface at a  
 63 temperature of 186 K that has been treated by  
 64 exposure to oxygen. It is found that such a sample  
 65 will emit thermal positronium with an efficiency of  
 66 about 12% and a beam-Maxwellian velocity distri-  
 67 bution [5]

$$dN/dE_{\perp} dE_{\parallel} = N_0(k_B T)^{-2} \exp\{-(E_{\perp} + E_{\parallel})/k_B T\}, \quad (1)$$

69 where  $E_{\perp}$  and  $E_{\parallel}$  are the components of the posi-  
 70 tronium energy perpendicular and parallel to the  
 71 surface respectively,  $k_B$  is Boltzmann's constant  
 72 and  $T$  is the sample temperature. The total number  
 73 of cold positronium atoms emitted from the hori-  
 74 zontal line source is taken to be  $N_0 = 7 \times 10^5$ . The  
 75 number of atoms emitted in a nearly normal di-  
 76 rection to the surface within a solid angle  $d\Omega$  and  
 77 having a maximum perpendicular energy  $E_{\perp}$ , is for  
 78 small  $E_{\perp}$ ,

$$N(E_{\perp}, \theta) = \frac{1}{2} N_0 (E_{\perp}/k_B T)^2 d\Omega = (v/3 \text{ m/ms})^4 d\Omega \times 3.6 \text{ atoms per pulse}, \quad (2)$$

where the perpendicular component of the posi- 80  
 tronium velocity is  $v$  and the positronium kinetic 81  
 energy is 82

$$E_{\perp} = (v/3 \text{ m/ms})^2 \times 51.1 \mu\text{eV}. \quad (3)$$

In a 0.5-mm region near the Al target, the slow 84  
 positronium atoms are to be excited to the 2S-state 85  
 using first-order Doppler-free two photon excita- 86  
 tion [6]. These atoms would then live about 1.1  $\mu\text{s}$  87  
 and travel about 3 mm, during which time they 88  
 would be excited to a high Rydberg d state with 89  
 principle quantum number  $n \approx 25$  by first-order 90  
 Doppler-free resonant absorption of pairs of circu- 91  
 larly polarized infrared photons. With correctly 92  
 aligned cw laser beams, no momentum will be 93  
 imparted to the Ps by the two-photon absorption. 94  
 By appropriate curvature of the two-photon wave 95  
 front, the positronium could be slowed to increase 96  
 the density of its low energy component. The 97  
 positronium is then to be spun up to the maximum 98  
 orbital angular momentum state with  $L = n - 1$  by 99  
 circularly polarized microwave radiation. 100

## 101 3. Free fall

We allow the positronium atoms to fall for  $q$  102  
 times the lifetime of the Rydberg state of the atoms 103  
 which is (see Fig. 2 and Ref. [7]) 104




Fig. 2. Radiative lifetimes of positronium excited states.

$$(n/25)^{5.236} \times 2.25 \text{ ms.}$$

$$(4) \quad \Delta x = \frac{1}{2}gt^2 = q^2(n/25)^{10.47} \times 24.8 \text{ } \mu\text{m.} \quad (6)$$

106 The total Ps flight path is then

$$(n/25)^{5.236} (v/3 \text{ m/ms})q \times 6.75 \text{ m.} \quad (5)$$

Note that the deflection of the positronium due to emission of a photon from the transition  $n \rightarrow n - 1, L \rightarrow L - 1$  is

108 The gravitational free fall is

110  
111  
112

$$\begin{aligned}\Delta p/p &= (1/4)\alpha^2(c/v)n^{-3} \\ &= 8.52 \times 10^{-5}(25/n)^3(3 \text{ m/ms}/v),\end{aligned}\quad (7)$$

114 which is so great as to preclude the emission of  
115 even a single photon during the free fall.

#### 116 4. Van der Waals mirror

117 The cold Rydberg Ps atoms could be focused by  
118 quantum mechanical reflection at a grazing angle  $\theta$   
119 from an elliptical dielectric mirror subtending a  
120 half angle  $\theta$  and solid angle  $d\Omega \approx \frac{1}{4}\sin^2\theta$ . One  
121 possibility is to illuminate the dielectric by a totally  
122 internally reflected wave so that an evanescent  
123 wave is present in the vacuum just outside the  
124 mirror. Appropriate choice of the frequency and  
125 intensity of the radiation would result in reflection  
126 of the positronium. Reflection will also occur for  
127 an unilluminated dielectric when the perpendicular  
128 component of the Ps wavelength is sufficiently  
129 greater than the length scale of the Van der Waals  
130 potential, i.e. when [8]

$$|dk_{\perp}/dx| \geq k_{\perp}^2, \quad (8)$$

132 where  $\hbar^2 k_{\perp}^2/4m_e = E_0 - V(x)$  is the perpendicular  
133 component of the positronium center of mass ki-  
134 netic energy, and  $E_0 = E_T \sin^2\theta$  is its value at very  
135 large  $x$ . The Van der Waals potential  $V(x)$  for a  
136 Rydberg Ps atom, with principle quantum number  
137  $n$  and  $L = n - 1$ , in vacuum a distance  $x$  from a flat  
138 surface with a dielectric constant  $\varepsilon$  is approxi-  
139 mately given by

$$\begin{aligned}V(x) &= -(\varepsilon - 1)(\varepsilon + 1)^{-1}(1/6) \\ &\quad \times (n^4 + 3n^3/2 + n^2/2)e^2 a_B^2/x^3,\end{aligned}\quad (9)$$

141 provided  $x > 2n^2 a_B$ , the mean radius of the Ryd-  
142 berg Ps atom defined by the expectation value of  
143  $1/r$ . Here  $r$  is the radial separation of the electron  
144 and positron and  $a_B$  is the Bohr radius of the in-  
145 finite mass hydrogen atom. We have also assumed  
146 that orbital angular momentum state is the one for  
147 which  $m = L$ , with the axis of quantization normal  
148 to the surface. The inequality  $|dk_{\perp}/dx| \geq k_{\perp}^2$  can-  
149 not be satisfied for

$$\begin{aligned}E_0 &> (2^{15}/3^{10})Ry(n^4 + 3n^3/2 + n^2/2)^{-2} \\ &\quad \times (\varepsilon + 1)^2(\varepsilon - 1)^{-2} \\ &= (25/n)^8(\varepsilon + 1)^2(\varepsilon - 1)^{-2} \\ &\quad \times (1 + 3/2n + 1/2n^2)^{-2} \times 49.5 \text{ peV.}\end{aligned}\quad (10)$$

We thus require

$$\begin{aligned}E_T &< (0.03/\sin^2\theta)(25/n)^8(\varepsilon + 1)^2(\varepsilon - 1)^{-2} \\ &\quad \times (1 + 3/2n + 1/2n^2)^{-2} \times 55 \text{ neV.}\end{aligned}\quad (11)$$

151 For a material like solid glass with  $\varepsilon \approx 3$ , the  
152 necessary kinetic energy of the positronium atoms  
153 will be unattainably small without laser cooling  
154 and phase space compression of the positronium.  
155 For a mirror surface made from a very low density  
156 material such as aerogel or liquid He with  
157  $\varepsilon = 1.04$ , we then have  
158  
159

$$\begin{aligned}E_T &< (0.03/\sin^2\theta)^2(25/n)^8 \\ &\quad \times (1 + 3/2n + 1/2n^2)^{-2} \times 138 \text{ } \mu\text{eV,}\end{aligned}\quad (12)$$

well within the velocity limit originally decided  
upon, i.e. 3 m/ms: 161 162

$$\begin{aligned}(v/3 \text{ m/ms}) &< (0.03/\sin\theta)(25/n)^4 \\ &\quad \times (1 + 3/2n + 1/2n^2)^{-1} \times 1.64.\end{aligned}\quad (13)$$

#### 5. Count rate

The total count rate calculated from Eq. (2) is  
now 166 167

$$\begin{aligned}N &= (v/3 \text{ m/ms})^4(\sin\theta/0.03)^2 e^{-(q-1)}(\eta/0.01) \\ &\quad \times 3.0 \times 10^{-4} \text{ counts per second,}\end{aligned}\quad (14)$$

169 with  $\eta$  being the product of the net optical exci-  
170 tation probability, the probability of reflection  
171 from the elliptical mirror and the detection effi-  
172 ciency. The positron pulse repetition rate has been  
173 taken to be  $100 \text{ s}^{-1}$ . Given the careful optimization  
174 of  $\eta$ , this rate would be sufficient to allow for fo-  
175 cusing, alignment and systematic tests, most of  
176 which however would be done using the faster  
177 positronium atoms. Note that the trajectories of  
178 the fast positronium atoms defines the zero of

179 deflection. From Eq. (6), the magnitude of the  
180 largest deflection will be of order  $50 \mu\text{m}$  for  $q = 1.5$   
181 and  $n = 25$ . This should be readily measured using  
182 a channel plate detector having channels of di-  
183 ameter  $10 \mu\text{m}$ .

## 184 6. Extraneous influences

185 There are many perturbations that can disturb a  
186 measurement on a single atom falling under the  
187 influence of gravity [9]. The gravitational force on  
188 a positronium atom in the earth's field is

$$F_G = 5.1 \times 10^{-13} \text{ eV/cm.} \quad (15)$$

190 The image potential of a metal surface a distance  $x$   
191 from Rydberg Ps will produce a force

$$F_I = (Ry/a_B)(na_B/x)^4, \quad (16)$$

193 which will exceed the gravitational force for  
194  $x < 300 \mu\text{m}$ .

195 The polarizability of the Ps is approximately  
196 equal to  $r^3$ , and its potential energy in an electric  
197 field  $E$  is  $U = r^3 E^2$ . The patch potential variations  
198 [10] a distance  $x$  from a surface having uniformly  
199 distributed patches of size  $\delta$  with work function  
200 variations  $\Delta\phi$  will be  $\Delta\phi (\delta/x)$  due to the averaging  
201 of the fluctuations. The electric fields will be of  
202 order  $E \approx \Delta\phi\delta/ex^2$ . The patch potential force will  
203 then be of order

$$\begin{aligned} F_P &= (\Delta\phi\delta/ex^2)^2 r^3/x \\ &= (\Delta\phi/0.5 \text{ eV})^2 (\delta/0.1 \mu\text{m})^2 (n/25)^6 \\ &\quad \times (300 \mu\text{m}/x)^5 \times 2 \times 10^{-14} \text{ eV/cm,} \end{aligned} \quad (17)$$

205 which will be negligible compared to  $F_G$  except in  
206 the vicinity of the mirror. How to maintain low  
207 stray fields near the mirror is an unsolved problem  
208 since it must have a low dielectric constant. Pos-  
209 sibly it could be made from aerogel with a thin  
210 coating of colloidal graphite.

211 The stray electric fields will cause Stark mixing  
212 and thus quenching and ionization of the Rydberg  
213 states. Also to be avoided are effectively time  
214 varying electric fields due to the motion of the Ps  
215 through spatially varying fields. Excitations with  
216  $\Delta n = 1$  for  $n = 25$  will occur at roughly  $80 \text{ GHz}$   
217 due to spatial variations of about  $3 \times 10^5 \text{ cm}^{-1}$

218 that are only present within a few hundred nm of a  
219 surface. Transitions with  $\Delta n = 0$  will require much  
220 slower variations and need to be examined closely.

221 Collisions with residual gas atoms in a 10-m  
222 flight path at  $10^{-11} \text{ Torr}$ , for which the density is  
223  $3 \times 10^5 \text{ molecules/cm}^3$ , would be just negligible for  
224 a cross-section of  $4\pi r^2 \approx 10^{-9} \text{ cm}^2$ . Magnetic fields  
225 will mix the orbital substates, introduce motional  
226 Stark quenching and also deflect the atoms.

227 Thermal radiation from the walls of the vacuum  
228 container can photoionize the positronium or in-  
229 duce transitions. Liquid He temperature has  $k_B T$   
230 just the same as the  $\Delta n = 1$  transition energy for  
231  $n = 25$ . The transition rates need to be calculated.

## 7. Conclusion

233 We can hardly claim at this point that a free fall  
234 measurement would be practical, given the possi-  
235 bility of unanticipated systematic effects and the  
236 difficulty of accounting for some of the known  
237 disturbances. However the experiment seems  
238 promising enough to warrant further calculations  
239 and preliminary experiments.

## References

- 241 [1] W.M. Fairbank, F.C. Witteborn, J.M.J. Madey, J.M.  
242 Lockhart, in: B. Bertotti (Ed.), Experimental Gravitation,  
243 Academic Press, NY, 1974, p. 310.  
244 [2] G. Chardin, J.M. Rax, Phys. Lett. B 282 (1992) 256;  
245 G. Chardin, Nucl. Phys. A 558 (1993) 447;  
246 G. Chardin et al., Phys. Rep. 241 (1994) 65.  
247 [3] A.P. Mills Jr., Hyperfine Interact. 44 (1988) 107 (Fig. 22).  
248 [4] T.J. Phillips, Hyperfine Interact. 109 (1997) 357.  
249 [5] A.P. Mills Jr., E.D. Shaw, M. Leventhal, R.J. Chichester,  
250 D.M. Zuckerman, Phys. Rev. B 44 (1991) 5791.  
251 [6] M.S. Fee, S. Chu, A.P. Mills Jr., R.J. Chichester, D.M.  
252 Zuckerman, E.D. Shaw, K. Danzmann, Phys. Rev. A 48  
253 (1993) 192.  
254 [7] H.A. Bethe, E.E. Salpeter, Quantum Mechanics of One-  
255 and Two-electron Atoms, Springer-Verlag, Berlin, 1957 (p.  
256 266, Table 15).  
257 [8] T. Martin, R. Bruinsma, P.M. Platzman, Phys. Rep. 223  
258 (1993) 135;  
259 C. Carraro, M.W. Cole, Progr. Surf. Sci. 57 (1998) 61.  
260 [9] T.W. Darling, F. Rossi, G.I. Opat, G.F. Moorhead, Rev.  
261 Mod. Phys. 64 (1992) 237.  
262 [10] J.B. Camp, T.W. Darling, R.E. Brown, J. Appl. Phys. 71  
263 (1992) 783, 69 (1991) 7126-7129.